

## BACKGROUND

Signal transducers and activators of transcription (STATs) were first discovered in protein complexes bound to the promoters of interferon-inducible genes. Seven STATs have now been cloned: STAT1, STAT2, STAT3, STAT4, STAT5A, STAT5B, and STAT6. These transcription factors participate in normal cellular events, such as differentiation, proliferation, cell survival, apoptosis, and angiogenesis following cytokine, growth factor, and hormone signaling.<sup>1</sup> STATs are activated by tyrosine phosphorylation, which is normally a transient and tightly regulates process. The seven mammalian STATs bear a conserved tyrosine residue near the C-terminus that is phosphorylated by JAKs. This phosphotyrosine permits the dimerization of STATs through interaction with a conserved SH2 domain. Phosphorylated STATs enter the nucleus by a mechanism that is dependent on importin-5 (also called nucleoprotein interactor 1) and the Ran nuclear import pathway. Once in the nucleus, dimerized STATs bind specific regulatory sequences to activate or repress transcription of target genes. Thus the JAK/STAT cascade provides a direct mechanism to translate an extracellular signal into a transcriptional response.<sup>2</sup> In addition, serine phosphorylation of STATs may be required for the maximal induction of Stat-mediated gene activation.<sup>3</sup> STATs can be divided into two groups according to their specific functions. One is made up of STAT2, STAT4, and STAT6, which are activated by a small number of cytokines and play a distinct role in the development of T-cells and in IFN signaling. The other group includes STAT1, STAT3, and STAT5, activated in different tissues by means of a series of ligands and involved in IFN signaling, development of the mammary gland, response to GH, and embryogenesis. This latter group of STATs plays an important role in controlling cell-cycle progression and apoptosis and thus contributes to oncogenesis.

Stat1 protein exists as a pair of isoforms, Stat1 $\alpha$  (91 kDa) and the splice variant Stat1 $\beta$  (84 kDa). Although an increased expression of STAT1 has been observed in many human neoplasias, this molecule can be considered a potential tumor suppressor, since it plays an important role in growth arrest and in promoting apoptosis.<sup>4</sup> On the other hand, STAT3 and 5 are considered as oncogenes, since they bring about the activation of cyclin D1, c-Myc, and bcl-xl expression, and are involved in promoting cell-cycle progression, cellular transformation, and in preventing apoptosis.<sup>5</sup>

Stat5B mediates the signal transduction triggered by various cell ligands, such as IL2, IL4, CSF1, and different growth hormones. It has been shown to be involved in diverse biological processes, such as TCR signaling, apoptosis, adult mammary gland development, and sexual dimorphism of liver gene expression.<sup>6</sup> Stat5B gene was found to fuse to

retinoic acid receptor-alpha (RARA) gene in a small subset of acute promyelocytic leukemias (APLL). The dysregulation of the signaling pathways mediated by this protein may be the cause of the APLL.

### References:

1. Artavanis-Tsakonas S et al.: Science 284: 770-776, 1999.
2. Oswald F et al.: Mol. Cell. Biol. 21: 7761-74, 2001.
3. Laky K & Fowlkes BJ: Curr. Opin. Immunol. 20: 197-202, 2008.
4. Chen, S. & Schreiber, S.: Cancer Res. 69(24 Suppl):Abstract nr 65, 2009
5. Calò, V. et al: J. Cell. Physiol. 197:157-68, 2003
6. Teglund, S. et al: Cell 93:841-50, 1998

## TECHNICAL INFORMATION

### Source:

Stat5B antibody is a mouse monoclonal antibody raised against recombinant human Stat5B fragments expressed in *E. coli*.

### Specificity and Sensitivity:

This antibody detects endogenous Stat5B proteins without cross-reactivity with other family members.

**Storage Buffer:** PBS and 30% glycerol

### Storage:


Store at -20°C for at least one year. Store at 4°C for frequent use. Avoid repeated freeze-thaw cycles.

## APPLICATIONS

Application:	*Dilution:
WB	1:1000
IP	1:50
IHC	n/d
ICC	n/d
FACS	n/d
*Optimal dilutions must be determined by end user.	



### QUALITY CONTROL DATA



Western Blot detection of Stat5B proteins in various cell lysates using Stat5B Antibody.

