

RECOMBINANT PROTEINS Mouse Resistin-Like Molecule-Gamma, Animal-Free **CAT. NO.** RP2089AF-25 RP2089AF-100 RP2089AF-1000

BACKGROUND:

Resistin-like molecule-gamma (RELM- γ) is a member of the RELM family of secreted proteins containing C-terminal cysteines. The RELM family consists of Resistin (FIZZ3), RELM-a (FIZZ1), RELM- β (FIZZ2), and RELM- γ (FIZZ4). RELM- γ is secreted by peripheral blood granulocytes, bone marrow, spleen, intestine, and lung. RELM- γ functions to promote and regulate promyelocytic differentiation, in addition to regulating nutrient-associated insulin sensitivity in the intestinal tract. Rodents secrete all four RELM family members, whereas Resistin and RELM- β are the only RELM family members found in humans.

Recombinant mouse Resistin-like molecule-gamma is a non-glycosylated protein dimer, containing 89/178 amino acids and having a molecular mass of 9.4/18.9 kDa.

Cat. No.: RP2089AF

AA Sequence:

MEGTLESIVE FSCTSITASG GSWDIRDGNT KKVKELLANR D RLASCPSGMT V CHCQCSTMDW A

DDCPSTVTKT VTGCACGYGC ATARCCQLA

TECHNICAL INFO

SIZE

25 μg 100 μg

1000 uq

Source:

E. coli

Physical Appearance:

Sterile Filtered white lyophilized (freeze-dried) powder.

Formulation:

10 mM sodium phosphate, pH 7.5

Stability:

Lyophilized product is very stable at -20° C. Reconstituted material should be aliquoted and frozen at -20° C. It is recommended that a carrier protein (0.1% HSA or BSA) is added for long term storage.

Reconstitution:

Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/mL, which can be further diluted into other aqueous solutions.

Protein Content and Purity determined by:

- UV spectroscopy at 280 nm.
- RP-HPLC calibrated against a known standard.
- Quantitation against a known standard via reducing and non-reducing SDS-PAGE gels.

Endotoxin Level:

Endotoxin level, as measured by LAL analysis, is <0.01ng/ug or <0.1EU/ug.

Animal Component-Free

This product is produced with no animal derived raw products. All processing and handling employs animal free equipment and animal free protocols.

