

BACKGROUND

c-Abl is a non-receptor protein-tyrosine kinase related to the Src family. The N-terminal region of c-Abl is structurally similar to Src and includes Src homology regions 3 and 2 (SH3 and SH2) and the catalytic domain; however, Abl differs from all Src family members by the presence of a large (90 kDa) C-terminal domain. Several functional motifs have been defined in the Abl C-terminus, including a nuclear localization signal (NLS), DNA binding domain, actin binding domain, sites for phosphorylation by protein kinase C and cdc-2 kinase, and binding sites for the adapter proteins Crk, Grb-2 and Nck.¹ c-Abl has been implicated in many cellular processes including differentiation, division, adhesion, death, and stress response. c-Abl becomes activated in response to numerous extra- and intra-cellular stimuli.² c-Abl is localized in the nucleus and the cytoplasm, where it plays distinct roles. The effects of c-Abl are mediated by multiple protein-protein and protein-DNA interactions and its tyrosine kinase domain.³ Its oncogenic counterpart, the Bcr-Abl fusion protein, causes particular human leukaemias. Use of targeted therapy that inhibits BCR-ABL kinase activity by Gleevec leads to hematologic and cytogenetic responses in affected individuals.⁴

References:

1. Hantschel, O. & Superti-Furga, G.: Nat Rev Mol Cell Biol. 5:33-44, 2004.
2. Sirvent, A. et al: Biol Cell 100:617-631, 2008.
3. Wang, J.Y.: Nat Cell Biol. 6:3-7, 2004.
4. Curran, M. P. et al: Biodrugs, 18:207-210, 2004.

TECHNICAL INFORMATION

Source:

Phospho-c-Abl (Tyr412) antibody is a rabbit polyclonal antibody raised against the epitope surrounding and including Tyr412 of the human c-Abl sequence.

Specificity and Sensitivity:

This affinity purified antibody detects endogenous phospho-human, mouse and rat c-Abl proteins.

Storage Buffer: PBS and 30% glycerol

Storage:


Store at -20°C for at least one year. Store at 4°C for frequent use. Avoid repeated freeze-thaw cycles.

APPLICATIONS

Application:	*Dilution:
WB	1:1000
IP	n/d
IHC (Paraffin)	n/d
ICC	n/d
FACS	n/d

**Optimal dilutions must be determined by end user.*

QUALITY CONTROL DATA



K562 cells were treated with Gleevec and subjected to Western Blot analysis using phospho-c-Abl (Tyr412) rabbit polyclonal antibody (**top**), or c-Abl antibody (**bottom**).

